

# On the Form of Odd Perfect Gaussian Integers

Matthew Ward

May 14, 2008

## 1 Introduction

Let  $\mathbb{Z}[i] = \{a + bi : a, b \in \mathbb{Z}\}$  be the ring of Gaussian integers. All Gaussian integers will be represented by Greek letters and rational integers by ordinary Latin letters. Primes will be denoted by  $\pi$  and  $p$  respectively. Units will be denoted by  $\varepsilon = \pm 1, \pm i$  and 1 respectively. In 1961 Robert Spira [3] defined the sum-of-divisors function on  $\mathbb{Z}[i]$  as follows. Let  $\eta = \varepsilon \prod \pi_i^{k_i}$  be a Gaussian integer. This representation is unique in that we will choose our unit  $\varepsilon$  such that each  $\pi_i$  is in the first quadrant ( $\text{Re}(\pi_i) > 0$  and  $\text{Im}(\pi_i) \geq 0$ ). Spira defined the sum-of-divisors function  $\sigma$  as

$$\sigma(\eta) = \prod \frac{\pi_i^{k_i+1} - 1}{\pi_i - 1}$$

This can be seen to match precisely with the rational form of the sum-of-divisors exactly when the rational primes coincide with the gaussian primes (when  $p \equiv 3 \pmod{4}$ ).

A Gaussian integer  $\eta$  is considered *even* if and only if  $1 + i$  divides  $\eta$ . It is easy to see that  $1 + i$  divides  $a + bi$  if and only if  $a$  and  $b$  have the same parity, i.e.,  $a \equiv b \pmod{2}$ . It follows that the usual parity rules for addition for  $\mathbb{Z}$  hold in  $\mathbb{Z}[i]$ : The sum of two Gaussian integers of the same parity is even, and of opposite parity, odd. Also, since  $a \equiv a^2 \pmod{2}$  for any rational integer  $a$ , it follows that a Gaussian integer  $a + bi$  is even if and only if its norm,  $N(a + bi) = a^2 + b^2$ , is an even rational integer.

Finally, we can define *perfect numbers* in the natural way. A Gaussian integer  $\eta$  is *perfect* if  $\sigma(\eta) = (1 + i)\eta$ . This is not the only notion we can work with, though. The Gaussian integer  $\eta$  is *norm-perfect* if  $N(\sigma(\eta)) = N(1 + i)N(\eta) = 2N(\eta)$ . Every perfect number is norm-perfect, so often it is easier to work with the norm perfect concept. Wayne McDaniel [1] proved a theorem for Gaussian integers analogous to Euclid's and Euler's characterization of even perfect positive integers. We will prove a theorem for Gaussian integers analogous to Euler's partial characterization of odd perfect positive integers, which is that

---

<sup>0</sup>Research for this paper was completed at the Summer 2007 Research Experience for Undergraduates at Auburn University, under the auspices of NSF grant no. 03533723.

any such integer must be of the form  $p^j m^2$  where  $\gcd(p, m) = 1$  and  $p \equiv j \equiv 1 \pmod{4}$ .

## 2 The Form of Odd Perfect Numbers

**Lemma 2.1.** *If  $\pi$  is an odd prime, then  $\sigma(\pi^m)$  is even if and only if  $m$  is odd.*

*Proof.* We first claim that for any Gaussian integers  $\eta_j$  we have  $N\left(\sum_j \eta_j\right) \equiv \sum_j N(\eta_j) \pmod{2}$ . This follows from basic parity rules. We have  $N\left(\sum_j \eta_j\right) \equiv 0 \pmod{2}$  if and only if  $\sum_j \eta_j$  is even in the Gaussian sense, but this happens if and only if there are an even number of odd  $\eta_j$ 's. This means that there are an even number of  $N(\eta_j) \equiv 1 \pmod{2}$ . Therefore, we have that  $N\left(\sum_j \eta_j\right) \equiv 0 \pmod{2}$  if and only if  $\sum_j N(\eta_j) \equiv 0 \pmod{2}$ .

Suppose  $\pi$  is an odd prime. Now examine  $N(\sigma(\pi^m)) = N(1 + \pi + \cdots + \pi^m) \equiv N(1) + N(\pi) + \cdots + N(\pi^m) \pmod{2}$ . We clearly have that this sum is congruent to 0 mod 2 if and only if  $m$  is odd since each term in the sum is odd.  $\square$

**Theorem 2.2.** *If  $\alpha$  is an odd norm-perfect Gaussian integer, then  $\alpha = \pi^k \gamma^2$  where  $k$  is an odd rational integer and  $\gcd(\pi, \gamma) = 1$ .*

*Proof.* Let  $\alpha$  be an odd norm-perfect Gaussian integer. Then  $\alpha = \prod_{i=1}^n \pi_i^{k_i}$ , where no  $\pi_j$  is associate to  $(1+i)$ . Since  $\alpha$  is norm-perfect we have  $N(\sigma(\alpha)) = 2N(\alpha)$ , but since  $N(\alpha)$  is odd and the sum of two squares we have that  $N(\alpha) \equiv 1 \pmod{4}$ , so  $N(\sigma(\alpha)) \equiv 2N(\alpha) \equiv 2 \pmod{4}$ .

Examine

$$\begin{aligned} N(\sigma(\alpha)) &= \prod_{i=1}^n N(\sigma(\pi_i^{k_i})) \\ &= N(\sigma(\pi_1^{k_1})) \cdots N(\sigma(\pi_n^{k_n})) \end{aligned}$$

Without loss of generality we can suppose that  $N(\sigma(\pi_1^{k_1})) \equiv 2 \pmod{4}$  and all the other terms in the product above are congruent to 1 mod 4. This is because no term can be 0 mod 4 or else the whole product would be 0 mod 4, and no term can be 3 mod 4 since the norm is the sum of two squares. From Lemma 2.1 we know that  $k_1$  is odd and that each  $k_i$  is even for all  $1 < i \leq n$ .

The form of an odd norm-perfect Gaussian integer then must have one prime to an odd power and the rest of the factorization are squares. Thus the form is  $\alpha = \pi^k \gamma^2$  where  $k$  is odd and  $\gcd(\pi, \gamma) = 1$ .  $\square$

**Corollary 2.3.** *If  $\alpha$  is an odd perfect Gaussian integer, then  $\alpha = \pi^k \gamma^2$  where  $k$  is an odd rational integer and  $\gcd(\pi, \gamma) = 1$ .*

*Proof.* Every perfect Gaussian integer is norm-perfect.  $\square$

### 3 Some Pesky Counterexamples

As noted above, in the rational case, Euler proved that every odd perfect number has the form  $p^j b^2$  where  $p$  is a prime,  $\gcd(p, b) = 1$ ,  $p \equiv 1 \pmod{4}$ , and  $j \equiv 1 \pmod{4}$ . In our form,  $\pi_1^k \gamma^2$ , we get that  $N(\pi_1) \equiv 1 \pmod{4}$  for free since the norm is a sum of two squares. However, we cannot show that  $k \equiv 1 \pmod{4}$ .

It is still unknown whether or not there are any odd perfect Gaussian integers, but there are odd norm-perfect numbers. In fact the smallest is a rather disturbing example:  $2 + i$ . It turns out that this is not only odd, but a prime. Have no fear, we shall show that this is the only such example. Let  $\pi$  be a Gaussian prime. If  $\pi$  is norm-perfect then  $N(\pi + 1) = 2N(\pi)$ ; if  $\pi = a + bi$  then this equation becomes

$$(a + 1)^2 + b^2 = 2(a^2 + b^2)$$

which is equivalent to  $(a - 1)^2 + b^2 = 2$ . The only 4 integer solutions to this are  $(2, -1)$ ,  $(0, -1)$ ,  $(2, 1)$ ,  $(0, 1)$ . Therefore in the Gaussian integers  $2 + i$  and  $2 - i$  are the only primes that are norm-perfect.

### 4 Acknowledgments

I would like to thank the NSF and Auburn University for providing the REU and facilities for this research. I would like to acknowledge Pete Johnson of Auburn University for advising me and providing comments on this paper. I also acknowledge Jeff Ward for introducing me to the sum-of-divisors and allowing me to bounce ideas off of him all summer.

### References

- [1] Wayne L. McDaniel. Perfect gaussian integers. *Acta Arithmetica*, 25:137–144, 1974.
- [2] Joseph B. Muskat. On divisors of odd perfect numbers. *Mathematics of Computation*, 20(93):141–144, 1966.
- [3] Robert Spira. The complex sum of divisors. *American Mathematics Monthly*, 68(2):120–124, 1961.